





Giving back control: Improving the maternity care experience of survivors of sexual violence


Katherine Talbot and Wendy Randall
Oxford University Hospitals
Ellie Bennett
Oxford Sexual Abuse and Rape Crisis Centre
15th March 2017


My body is the evidence
The spoils of his crime.
It is tainted and dirty –
Never to be mine.

I cringe and I shudder,
The memories still hurt,
Of the baby – the child –
The woman of dirt.









Oxford Academic Health Science Network

Table 3. Matemal daily cigarette smoking and illicit drug use at some point during first subsequent pregnancy by violence severity, age and

| | Smoking of ≥10 cigarettes | | | | Illicit drug use | | | |
|--|---------------------------|------|------|-----------|------------------|-----|-----------------|------------|
| | na | % | RRb | 95% CI | na | % | RR ^b | 95% CI |
| Unexposed | 1489 | 4.4 | ref. | | 1468 | 0.4 | ref. | |
| Severity of the exposure | | | | | | | | |
| First pregnancy after any sexual violence | 540 | 15.0 | 3.55 | 2.46-5.12 | 523 | 3.8 | 6.27 | 2.13-18.43 |
| First pregnancy after penetrative sexual violence | 387 | 15.8 | 3.71 | 2.51–5.49 | 376 | 4.0 | 7.33 | 2.25–23.87 |
| Age at time of first attendance ^c | | | | | | | | |
| <20 years | 311 | 12.2 | 2.17 | 1.33-3.54 | 302 | 3.6 | 2.46 | 0.95-6.39 |
| ≥20 years | 227 | 18.9 | 4.04 | 2.81-5.83 | 219 | 4.1 | 10.89 | 3.93-30.13 |
| Missing | 2 | | | | 2 | | | |
| Time from assault to first delivery ^c | | | | | | | | |
| <3 years | 217 | 20.3 | 3.55 | 2.27-5.54 | 209 | 5.7 | 9.52 | 2.62-34.64 |
| 3–6 years | 190 | 15.3 | 3.54 | 2.18-5.74 | 184 | 3.3 | 4.82 | 1.26-18.47 |
| ≥6 years | 127 | 6.3 | 2.19 | 0.98-4.89 | 125 | 1.6 | 2.20 | 0.48-10.12 |
| Missing | 6 | | | | 5 | | | |

^aNumber of women with available information and the proportion with each risk factor.

Risk factors and health during pregnancy among women previously exposed to sexual violence

Acta Obstet Gynecol Scand 2014; 93:351-358.

Odds ratios (and 95% confidence intervals) for thoughts and wishes about birth for previoussexual violence, stratified by parity.


| Nullipara | No violence $n=35,570$ | Previous sexual violence $n = 7280$ | | | | |
|-------------------------------|------------------------|-------------------------------------|-----------------------------------|-------------------------------|--|--|
| | % | % | OR | AOR* | | |
| Fear of childbirth | 16.4 | 21.8 | 1.4 (2.3-2.5) | 1.2 (1.1-1.3) | | |
| Hope for a a pain-free birth | 7.4 | 8.8 | 1.2 (1.1-1.3) | 1.0 (0.9-1.1) | | |
| Epidural regardless | 9.1 | 10.8 | 1.2 (1.1-1.3) | 1.0 (0.9-1.1) | | |
| Desire for caesarean section | 3.2 | 4.6 | 1.5 (1.3-1.7) | $(1.2 (1.0-1.4)^{\dagger})$ | | |
| Worries about infant's health | 11.2 | 16.0 | 1.5 (1.4-1.6) | 1.2 (1.1-1.3) | | |
| Looking forward to infant | 97.1 | 96.0 | 0.7 (0.6-0.8) | 0.8 (0.7 (0.9) | | |
| Multipara | No violence n=28,622 | | Previous sexual violence $n=6394$ | | | |
| | % | % | OR | AOR* | | |
| Fear of childbirth | 20.6 | 25.4 | 1.3 (1.2-1.4) | (1.1 (1.0-1.2) [†]) | | |
| Hope for a pain-free birth | 8.6 | 9.6 | 1.1 (1.9-1.2) | 1.0 (0.9=1.1) | | |
| Epidural regardless | 10.7 | 12.6 | 1.2 (1.1-1.3) | 11 (1.0-12) | | |
| Desire for caesarean section | 6.2 | 7.4 | 1.2 (1.1-1.4) | (1.0 (0.9-1.1)) | | |
| Worries about infant's health | 10.7 | 14.1 | 1.4 (1.3-1.5) | 1.1 (1.0-1.2) | | |
| Looking forward to infant | 97.2 | 96.0 | 0.7 (0.6-0.8) | 0.8 (0.7-0.9) | | |


Lifetime sexual violence and childbirth expectations – A Norwegian population based cohort study Midwifery 36(2016)14-20


^DRelative risks adjusted for age, parity and year of delivery.

^CSome women attended the Rape Trauma Service more than once. Age categorization is according to age at first attendance. Time between assault and delivery was calculated from the most recent assault to date of delivery.

 ^{§ =} Additionally controlled for a previous caesarean section.
 * Controlled for: Age, education, civil status, pre pregnancy risk factors and mental distress.













- Increasing understanding of sexual violence
- Understand how sexual violence can impact on pregnancy
- Build confidence in responding to women's fears and concerns about pregnancy arising from sexual violence
- Consider how we can empower women to make positive choices about their pregnancy












Next time ... More time on ... everything! Interview with a survivor Legal Aspects Network Development



The Future

- > Training available to everyone
- > Survivors of sexual violence feel empowered
- Thames Valley is know for its care, not its crime

